

A STUDY OF MATHEMATICAL FORMULATION ON INITIAL-BOUNDARY VALUE PROBLEMS FOR FIRST ORDER PDES IN UNBOUNDED DOMAINS

Neha¹ and Jakhar, Manjeet Singh²

¹Research Scholar, NIILM University, Kaithal, Haryana

²Associate Professor, Department of Mathematics, NIILM University, Kaithal, Haryana

DOI: <https://doi.org/10.70388/ijabs250113>

Received on Sep 10, 2024

Accepted on Nov 9, 2024

Published on Jan 10, 2025

This article is licensed under a license [Commons Attribution-NonCommercial-NoDerivatives 4.0 International Public License \(CC BY-NC-ND 4.0\)](#)

Abstract

At some point in the solution domain of a differential equation, there occurs an initial value issue (also known as the Cauchy problem) in the company of a stated value known as starting situation of arbitrary derivative or function. which is a topic within partial differential equations. In physics and other fields, addressing an initial value issue is a common part of modelling a system. Some fields of physics, such as quantum mechanics and relativity, include boundary value difficulties, as any physical differential equation will. Boundary value difficulties, such as determining the normal nodes, are common in the wave equation. Similarities between boundary value and start value issues may be found. In a boundary value problem, all of the conditions are illustrated at same value of the independent variable, while in an initial value problem, all of the criteria are provided at various independent variable values. The initial value is termed (the value is at the lower boundary of the domain thus, the term initial value). An initial value issue and a boundary value problem must be well-posed in order to be relevant in applications. This indicates that given the input to the issue, there is a single solution that is constantly dependent on the input.

Keywords: Mathematical Formulation, Initial-Boundary, Value Problems, First Order PDES, Unbounded Domains, differential equation

Introduction

First-order hyperbolic equations with two independent variables x and t are solved using a finite difference method. Typically, the first variable is space, while the second is time. For hyperbolic equations, a fundamental difficulty is constructing finite difference schemes that are both stable and do not damp out of the solution. Initial and initial-boundary value approximation algorithms are developed in this suggested study.

Motivation

First-order hyperbolic partial differential equations have a rich literature. It took a long time and a lot of work to create parameteric estimation techniques for Initial value and Initial-boundary value issues/problems and initial-boundary value problems. Gottlieb et al (1987), Bo (1998), and Coulombel (2009) investigated the performance of finite - element strategies for 1st first order hyperbolic initial-boundary value problems using vectors value functions having $L_2(\mathbb{R}^+, \mathbb{R}^N)$. Semi-distributed approximations to the starting and boundary value issue were addressed by Warming and Beam in 1988.

$$U_t = aU_x, \quad 0 \leq x \leq A, t \geq 0,$$

$$U(x, 0) = u(x), \quad 0 \leq x \leq A,$$

$$U(A, t) = v(t), \quad t \geq 0,$$

(1)

Wherein $L_2[0, A]$ is " $a > 0$ " and " $v(t) = 0$ ". Using wavelets, Sekino and Hamada in 2008 derived a numerical solution to the Advection problem $u_t + (a(x)u)_x = 0$. Despres and Teng in 2009 and 2010 respectively developed finite-element techniques for the initial value problem. $U_t + aU_x = 0, x \in \mathbb{R}, t \in \mathbb{R}^+$,

$$U(x, 0) = u_0(x), \quad x \in \mathbb{R} \dots \quad (2)$$

For constrained initial functions u_0 with discontinuous starting values. Motivated by the development of numerical methods to start initial-boundary value problems, these works are published.

Problem Formulation

Initial Value Issue (IVP) on an infinite interval is the initial model problem in this suggested study.

$$U_t + a(x)U_x(x), x \in \mathbb{R}^+, t \in \mathbb{R}^+,$$

$$U(x,0) = u(x), x \in \mathbb{R}^+, \dots (3)$$

$A(x)$ is greater than or equal to zero for all nonnegative values of x between \mathbb{R}^+ and $C(\mathbb{R}^+)$, which is known as the initial condition. Wave propagation in homogeneous mediums is modelled by Equation (3).

IBVP (Initial-Boundary Value Issue) is an IB model problem that may be described as

$$U_t = -a U_x, x \in [0,1], t \in \mathbb{R}^+,$$

$$U(0, t) = v(t), t \in \mathbb{R}^+. \quad (4)$$

Assuming “ $a > 0$ ” & specification of the boundary condition $v(t)$ at $x = 0$. There are no errors in this boundary condition since the information is flowing from left to right and the compatibility criterion is met (0).

With IVP (3), there is no boundary condition, and hence, no IVP (4). Many scenarios need this latter requirement.

For the issues IVP (3) and IBVP (4), the semigroup theory was heavily used in order to construct a completely discrete convergent numerical method. To solve the initial-boundary value issue, semigroup theory offers an elegant solution.

PRELIMINARIES

Theorem of Pazy (1983) was the primary tool employed in this study, and this section provides basic definitions and a specific instance.

Theorem 1. Assume X is a Banach space consisting of standard $\| \cdot \|$. Assuming, X has denser $D(A)$ such that $A: D(A) \rightarrow X$ is a continuous (linear) projection. Further, λ is there, with

$\Re(\lambda) > 0$ wherein the range “ $\lambda I - A$ ” is dense in X . Let X_n be the Banach spaces consisting of standards $\| \cdot \|_n$. Furthermore, there are bounded linear operators, “ $P_n: X \rightarrow X_n$ and $E_n: X_n \rightarrow X$ ” for every $n \geq 1$ such that

- (i) $\|P_n\| \leq C_1, \|E_n\| \leq C_2$, with C_1 and C_2 represented as constants that are not dependent on n .
- (ii) $\|P_n x\|_n \rightarrow \|x\|$ as $n \rightarrow \infty$ for every $x \in X$.
- (iii) $\|E_n P_n x - x\| \rightarrow 0$ as $n \rightarrow \infty$ for every $x \in X$.
- (iv) $P_n E_n = I_n$ (I_n : identity operator on X_n)

Define $F(\tau_n)$ as a series of constrained linear operations spanning X_n to X_n fulfilling $\|F(\tau_n)^k\| \leq 1$. (5)

In addition to this, the constrained linear maps

$$A_n = \rho_n^{-1}(F(\rho_n) - I)$$

have the property that

$$D(A) = \{x \in X : E_n A_n P_n x \text{ converges}\}$$

and that

$$\lim_{n \rightarrow \infty} E_n A_n P_n x = A_x \quad (6)$$

for all $x \in D(A)$. A 's closure (\bar{A}) is therefore the indefinite constructor of contracting mappings “ $S(t)$ on X ”. Furthermore, if $k_n \tau_n \rightarrow t$ as $n \rightarrow \infty$, then

$$\lim_{n \rightarrow \infty} \|F((\tau_n)^{k_n})P_n x - P_n S(t)_x\|_n = 0$$

It is permissible to refer to a generalized, solution as "solutions" in the following paragraphs.

If $\alpha = (\alpha_0, \alpha_1, \dots, \alpha_k)$, then let the notation $\alpha(i) = \alpha_i$

For $x \in \mathbb{R}, [x] = \sup \{n \in \mathbb{Z} : n \leq x\}$

Theorem 2 Theorem (Hille-Yosida Theorem). An infinitesimal generator of the C_0 semigroup of contractions $T(t)$, $t \geq 0$ can only be generated by a linear (unbounded) operator A ,

- (i) $D(A) = X$, since A is closed
- (ii) \mathbb{R}^+ is present in $\overline{\rho(A)}$, the resolvent set of A . Further, The resolvent set $\rho(A)$ of A for every $\lambda > 0$, $\|R(\lambda; A)\| \leq \frac{1}{\lambda}$ (7)

IVP AND IBVP EXACT SOLUTIONS

For the start and initial-boundary value problems covered in this study, this section provides the precise solution.

IVP exact solution

The solution to eq (3) is provided by:

$$u(x, t) = u(\beta^{-1}(t + \beta(x))),$$

$$\text{Here, } \beta(x) = \int_0^x \frac{d\xi}{a(\xi)}$$

Using $u(x, t)$ data on a bounded domain, we were able to calculate the solution of (3) that was bound not compulsorily on a bounded domain, numerically. This conclusion is made possible by the following theorem.

Theorem 1. Assuming “ $\in C[0, \infty)$ ” and “ $a(x) > 0$ ” for every $x \in \mathbb{R}^+$. Assume $M > 0$ and $T > 0$.

Define $a_M: [0, M] \rightarrow \mathbb{R}$ as

$$a_M(x) = a(x), 0 \leq x \leq (M - \frac{1}{M})$$

$$= a(M - \frac{1}{M}) \sqrt{M(M - x)}, M - \frac{1}{M} \leq x \leq M$$

Letting $f \in C[0, M]$. The problem’s solution is

$$\frac{\partial V}{\partial t} = a_M(x) \frac{\partial V}{\partial x}, 0 \leq t \leq T, 0 \leq x \leq M,$$

$$V(x, 0) = f(x), 0 \leq x \leq M,$$

$$V(M, t) = f(M) \dots (8)$$

Is existing and unique in nature. It is provided as

$$V(x, t) = f(\beta_M^{-1}[\min(t + \beta_M(x), \beta_M(M))]),$$

$$\text{here, } \beta_M(x) = \int_0^x \frac{d\xi}{\alpha(\xi)}, 0 \leq x \leq M - \frac{1}{M}$$

$$= \int_0^{M-\frac{1}{M}} \frac{d\xi}{\alpha(\xi)} + \int_{M-\frac{1}{M}}^x \frac{d\xi}{\alpha\left(\left(M-\frac{1}{M}\right)\sqrt{M(M-x)}\right)}, M - \frac{1}{M} \leq x \leq M$$

Further,

$$S_t f(x) = f([\min(t + \beta_M(x), \beta_M(M))])$$

creates a contracting subclass on $C[0, M]$ with the generator

$$D(A) = \{g \in C[0, M]: g' \in C[0, M] \text{ and } \lim_{x \rightarrow M} a_M(x) g'(x) = 0\}$$

and

$$Ag(x) = a_M(x) g'(x), x \in [0, M],$$

$$Ag(M) = 0.$$

Furthermore, selecting “ $M > N$ ” in such a way that:

$$\sup_{t \in [0, T], x \in [0, N]} (t + \beta(x)) < \left(M - \frac{1}{M}\right),$$

$$V(x, t) = u(x, t), (x, t) \in [0, N] \times [0, T] \dots (9)$$

given “ $f \in C[0, M]$ ” as the constraint of u to $[0, M]$.

Proof. Defining $t \geq 0$, “ $T_t: [0, M] \rightarrow [0, M]$ ” as

$$T_t x = \beta_M^{-1}[\min(t + \beta_M(x), \beta_M(M))]$$

It is now simple to demonstrate that $T_{s+t} = T_s \circ T_t$.

$$\begin{aligned} T_s \circ T_t x &= \beta_M^{-1}[\min(s + \beta_M(T_t x), \beta_M(M))] \\ &= \beta_M^{-1}[\min(s + \beta_M(\beta_M^{-1}[\min(t + \beta_M(x), \beta_M(M))]), \beta_M(M))] \\ &= \beta_M^{-1}[\min([\min(s + t + \beta_M(x), s + \beta_M(M))], \beta_M(M))] \\ &= \beta_M^{-1}[\min(s + t + \beta_M(x), \beta_M(M))] \\ &= T_{s+t} x \end{aligned}$$

Additionally, because “ $S_t f(x) = f(T_t x)$ ”, it is simple to show that S_t is a subgraph.

Thus, according to the Hille-Yosida Theorem, Now, by Hille-Yosida Theorem, if S_t generator is B ,

Then,

$$\begin{aligned} (I - B)^{-1} h(x) &= \int_0^\infty e^{-t} S_t h(x) dt \\ &= \int_0^\infty e^{-t} h(\beta_M^{-1}[\min(t + \beta_M(x), \beta_M(M))]) dt \\ &= \int_0^{\beta_M(M) - \beta_M(x)} e^{-t} h(\beta_M^{-1}[(t + \beta_M(x), \beta_M(M))]) dt + \int_{\beta_M(M) - \beta_M(x)}^\infty e^{-t} h(N) dt \\ &= \int_0^M e^{\beta_M(M) - \beta_M(x)} \frac{h(y)}{a_M(y)} dy + h(N) e^{\beta_M(x) - \beta_M(M)} dt \end{aligned}$$

Here, $\beta_M^{-1}(t + \beta_M(x))$.

Considering the differential equation,

$$f(x) - a_M(x) f'(x) = h(x), x \in [0, M]$$

$$f(M) = h(M)$$

which is equivalent to

$$f(x) - a(x) f'(x) = h(x), x \in [0, M]$$

$$\lim_{x \rightarrow M} a(x) f'(x) = 0$$

Since, $\int_0^M \frac{ds}{a_M(s)} < \infty$, there exists unique solution for every “ $h \in X$ ” i.e., $f \in D(A)$ to the preceding differential equation that is provided as,

$$\begin{aligned} f(x) &= e^{\int_0^x \frac{ds}{a_M(s)}} \int_x^M \frac{h(y)}{a_M(y)} e^{-\int_0^y \frac{ds}{a_M(s)}} dy + h(M) e^{-\int_0^M \frac{ds}{a_M(s)}} + \int_0^x \frac{ds}{a_M(s)} \\ &= \int_x^M e^{\beta_M(x) - \beta_M(y)} \frac{h(y)}{a_M(y)} dy + h(M) e^{\beta_M(x) - \beta_M(M)} \end{aligned}$$

For the operators A and B, it can be demonstrated that $(I - A) 1 = (I - B) 1$.

“ $D(A) = D(B)$ ” and for any “ $g \in D(A)$ ”, “ $Bg = Ag$ ” may be deduced from this.

For $t \in [0, T]$ and $x \in [0, N]$, it is for sure a growing derivative/ function

$$x \leq \beta^{-1}(t + \beta(x)) < \left(M - \frac{1}{M}\right).$$

Hence,

$$\beta(\beta^{-1}(t + \beta(x))) = \beta_M(\beta_M^{-1}(t + \beta_M(x))).$$

This leads to the conclusion that “ $S_t f(x) = V(x, t) = u(x, t)$ ” for every “ $x \in [0, N]$ ” and “ $t \in [0, T]$ ”.

IBVP exact solution

Theorem 2. Assume “ $u \in C[0, 1]$ ” and “ $v \in C[0, \infty)$ ” in such a way that “ $u(0) = v(0)$ ”.

Defining $u_0(x) = u_0(x) - u_0(0)$. For “ U ” as a solution to (4), its also a solution to

$$\overline{U}_t = -a\overline{U}_x, \quad 0 \leq x \leq 1, t \geq 0$$

$$\overline{U}(x, 0) = -a\overline{U}_x, \quad 0 \leq x \leq 1 \quad (10)$$

and V is a solution to

$$V_x = \frac{-1}{a}V_t, \quad 0 \leq x \leq 1, t \geq 0$$

$$V(x, 0) = u_0(0), \quad 0 \leq x \leq 1,$$

$$V(0, t) = v(t), \quad t \geq 0 \dots \quad (11)$$

Then, “ $U(x, t) = U(x, t) + V(x, t)$ ”. Furthermore, for “ $T > 0$ ”, contracting subgroups s.t., are defined

“ $X \rightarrow X$ ” where “ $X = \{u \in C[0, 1]: u(0) = 0\}$ ” and “ $T(x): Y \rightarrow Y$ ”

Wherein, “ $Y = C[0, T]$ ” as

$$S_t u_0 = u_0(x - at), \quad at \leq x \leq 1$$

$$= 0, \quad 0 \leq x \leq at,$$

$$T_x w(t) = w(0), \quad 0 \leq x \leq x/a$$

$$= w\left(t - \frac{x}{a}\right), \quad t \geq x/a.$$

Then “ $U(x, t) = S_t u_0(x) + \sum x w(t)$ for all $(x, t) \in [0, 1] \times [0, T]$ ”.

W: represents constraint of “v to [0, T]”.

Furthermore, if A and B represent the St and Tx generators, respectively, then

“ $D(A) = \{g \in X: g' \in X\}$, $D(B) = \{g \in Y: g' \in Y \text{ and } g'(0) = 0\}$, $Ag = -ag'$ for all $g \in D(A)$ ” and “ $Bg = -1 a g'$ for all $g \in D(B)$ ”.

Convergent numerical technique for IVP and IBVP

M AND INITIAL-BOUNDARY VALUE PROBLEM

In this lesson, we'll go through how to solve the starting value and initial boundary value issues numerically convergently. It is possible to solve the initial value issue by posing it on a smaller bounding box, and then solving it on a larger bounding box with the same answer. An improved numerical solution to the modified issue is indistinguishable from the original answer on the smaller constrained region. Decomposing the initial-boundary value issue into two problems, each of which generates a semigroup, allows for the presentation of discrete semigroup approximations.

The IVP and a Convergent Numerical Scheme

$M > N$ and an initial value problem given on $[0, M]$ $[0, T]$ whose answer absolutely corresponds to the solution of (3) on $[0, N]$ $[0, T]$ for any subset of $[0, N]$ $[0, T]$ are possible for the initial value issue (3). \mathbb{R}^+ \mathbb{R}^+ \mathbb{R}^+ \mathbb{R}^+ \mathbb{R}^+ On $[0, M]$ $[0, T]$, create a finite difference scheme that converges to the solution of the problem provided in (3) on $[0, N]$ $[0, T]$.

This finding is made easier by the following theorem.

Theorem 1. Assume “ $X = C[0, M]$ ” & “A”. Let $X_n = \mathbb{R}^{n+1}$ whose elements are denoted as $\alpha = (\alpha_0, \alpha_1, \dots, \alpha_n)$. The supremum standard is applied to both X and X_n spaces. Finally, defining

$P_n: X \rightarrow X_n$ as $(P_n f)_i = f(iM/n)$, $i = 0, 1, \dots, n$.

$E_n: X_n \rightarrow X$ as

$E_n(\alpha)$ represents “piecewise linear function” having $E_n(\alpha)(iM/n) = \alpha_i$. Let

$$\tau_n = \frac{1}{2n(\sup_{x \in [0, M]} |a(x)|)}.$$

Defining operations “ $F(\tau_n): X_n \rightarrow X_n$ ” as

$$\begin{aligned} (F(\tau_n)\alpha)_i &= \left(1 - n\tau_n a_m \left(\frac{iM}{n}\right)\right) \alpha_i + \left(n\tau_n a_m \left(\frac{iM}{n}\right)\right) \alpha_{i+1}, i = 0, 1, \dots, n-1 \\ &= \alpha_n, i = n. \end{aligned}$$

Choosing $kn = \frac{t}{\tau_n}$, it can be shown that

$$\|\tau_n^{kn} P_n f - P_n S(t)f\|_n \rightarrow 0 \text{ as } n \rightarrow \infty \quad (12)$$

Particularly, fixing “ $t \in [0, T]$ ” & “ $x \in [0, N]$ ”,

$$\lim_{n \rightarrow \infty} F(\tau_n)^{kn} P_n f \left(\lfloor \frac{nx}{M} \rfloor\right) = u(x, t), \quad (13)$$

Where, where $u(x, t)$ represents a solution to (3).

Proof. Since P_n is clearly linear, $\|P_n\| = 1$. The fact that (ii) of Theorem 1 is true may be deduced with norms’ definitions from the definitions of the norms & constant continuance of component X . That $\|E_n\| = 1$ is simply obtained.

When you consider the definitions of E_n and P_n as well as the uniform continuation of the element X in Theorem 1, it becomes clear that (ii) is true. Formulating differential equation leads to the definition of $F(n)$ which is now simple. To illustrate, let's look at the $(iM/n, j_n)$ lattice functions for “ $i = 0, 1, 2, \dots$ ” and (“ $j = 0, 1, 2, \dots$ in the (x, t) plane”).

Assume $V(iM/n, j_n) = u_{i,j}$. Taking into account the $\sup_{x \in [0, M]} a_M(x) > 0$, τ_n is defined clearly. Take a look at the difference equation that corresponds to the differential equation in (8) is

$$\frac{u_{i,j+1} - u_{i,j}}{\tau_n} = a_m \left(\frac{iM}{n}\right) n(u_{i+1,j} - u_{i,j}), i = 0, 1, 2, \dots, n-1$$

$\frac{u_{n,j+1}-u_{n,j}}{\tau_n} = 0$, which can be simplified as

$$u_{i,j+1} = 1 - n\tau_n a_m \left(\frac{iM}{n}\right) u_{i,j} + n\tau_n a_m \left(\frac{iM}{n}\right) u_{i+1,j}, i = 0, 1, \dots, n-1$$

$$u_{n,j+1} = u_{n,j}.$$

It is possible to calculate all $u_{i,j}$ by using the preceding method for $u_{i,0}$. The formula for f_i is $f(iM/n) = f_i$.

Now,

$$\begin{aligned} & \|F(\tau_n)(\alpha)\|_n \\ &= \max \left(\max_{0 \leq i \leq n-1} \left| 1 - n\tau_n a_m \left(\frac{iM}{n}\right) \alpha_i + n\tau_n a_m \left(\frac{iM}{n}\right) \alpha_{i+1} \right|, |a_n| \right) \\ &= \max \left(\max_{0 \leq i \leq n-1} \left(1 - n\tau_n a_m \left(\frac{iM}{n}\right) \alpha_i + n\tau_n a_m \left(\frac{iM}{n}\right) \alpha_{i+1} \right), \max(|\alpha_i|, |\alpha_{i+1}|), |a_n| \right) \\ &= \max(|\alpha_0|, |\alpha_1|, \dots, |a_n|) \\ &= \|\alpha\|_n. \end{aligned}$$

Therefore, $\|F(\tau_n)\| \leq 1$, holding the theorem 1's stability condition (5).

For, $f \in D$,

$$\begin{aligned} & \left\| \tau_n^{-1} (F(\tau_n) - 1) P_n f - P_n a_m \left(\frac{iM}{n}\right) f \right\|_n \\ &= \sup_i \left| \frac{a_m \left(\frac{iM}{n}\right)}{n} \left(f \left(\frac{(i+1)M}{n}\right) - f \left(\frac{iM}{n}\right) \right) - a_m \left(\frac{iM}{n}\right) f \left(\frac{iM}{n}\right) \right| \dots \quad (14) \end{aligned}$$

Because " $f \in D$ ", " A_f " is consistently periodic on "[0, M]" the R.H.S of (14) goes to 0 as n tends to infinity ($n \rightarrow \infty$). As a result, Theorem 1 (3.6) is fulfilled.

Using Theorem 1, one must prove that for any $\epsilon > 0$, the range “ $I - A$ ” in X is dense. However, the range of $I - A$ is previously demonstrated to include all of X in Theorem 1. Theorem 1's formulation of $(I - A)^{-1}h$ may also be used.

CONCLUSION

Logic of partial functions as it is now understood is summarized and some directions for future research are suggested in this brief final study. This thesis was introduced by stating that partial functions have better logical and computational features than binary relations. This is worth repeating. This thesis's findings only serve to support this point of view. A first-order definition of an operation is one which may be defined according to the basic theorem. When it comes to these kinds of operations, it has already been proven that the representation classes are typically finitely axiomatisable and have equational theories of low complexity, the finite representation property is fulfilled and representability of finite algebras is easily decided. Moreover, similar observations seem to apply to functions with several locations. It's all the opposite of what happens in relationships. If we want to reason about programs represented by code written in any general-purpose (i.e., Turing-complete) language, we must be able to articulate some kind of unbounded iteration mechanism. However, translating in this manner yields results only if the signature contains anti-domain. Including anti-domain would be problematic if we were trying to describe partial recursive functions with no constraints, as finding the places where partial recursive functions are undefined is not an universal and effective computation that can be expressed in any programming language.

REFERENCES

1. Smith, H. An introduction to delay differential equations with applications to the life sciences, Text Book in Applied Mathematics, Vol. 57, First Edition, 2011.
2. Marcel Jackson and Tim Stokes, Modal restriction semigroups: towards an algebra of functions, International Journal of Algebra and Computation 21 (2011), no. 7, 1053–1095.
3. Ian Hodkinson, A construction of cylindric and polyadic algebras from atomic relation algebras, Algebra Universalis 68 (2012), no. 3, 257–285.
4. Robin Hirsch and Szabolcs Mikulas, Axiomatizability of representable domain algebras, The Journal of Logic and Algebraic Programming 80 (2011), no. 2, 75–91.

5. Robin Hirsch and Brett McLean, Disjoint-union partial algebras, *Logical Methods in Computer Science* 13 (2017), no. 2:10, 1–31.
6. Robin Hirsch, Marcel Jackson, and Szabolcs Mikulas, The algebra of functions with antidomain and range, *Journal of Pure and Applied Algebra* 220 (2016), no. 6, 2214–2239.
7. Robin Hirsch and Marcel Jackson, Undecidability of representability as binary relations, *The Journal of Symbolic Logic* 77 (2012), no. 4, 1211–1244.
8. Robert Goldblatt and Marcel Jackson, Well-structured program equivalence is highly undecidable, *ACM Transactions on Computational Logic* 13 (2012), no. 3, 26:1–26:8.
9. Simon Foster, Georg Struth, and Tjark Weber, Automated engineering of relational and algebraic methods in Isabelle/HOL, *Proceedings of 12th Conference on Relational and Algebraic Methods in Computer Science (Harrie de Swart, ed.)*, Springer, 2011, pp. 52–67.
10. Robert Egrot and Robin Hirsch, Completely representable lattices, *Algebra Universalis* 67 (2012), no. 3, 205–217.
11. Wiesław A. Dudek and Valentin S. Trokhimenko, Subtraction menger algebras, *Semigroup Forum* 85 (2012), no. 1, 111–128.
12. Cristiano Calcagno, Dino Distefano, Jeremy Dubreil, Dominik Gabi, Pieter Hooimeijer, Martino Luca, Peter O’Hearn, Irene Papakonstantinou, Jim Purbrick, and Dulma Rodriguez, Moving fast with software verification, *Proceedings of 9th NASA Formal Methods Symposium*, Springer, 2015, pp. 3–11.